

Paediatric Card - Inner Slide

Cut along this line before laminating

0	2mth	5mth	1	2	3	4	5	6	7	8	9	10	11	12	13	14	CPR Rates																	
																		Age	Ratio	Cycles														
3.5	5	7	10	12	14	16	18	20	22	24	26	33	36	40	43	46	Newborn	3 : 1	40 cycles/1 min															
40	20	20	20	20	20	20	20	20	20	20	15	15	15	15	15	15	Infant/Children	- (1 Person)	30 : 2	5 cycles/2 mins														
60	50	50	35	35	35	35	35	35	35	35	25	25	25	25	25	25	- (2 Person)	15 : 2	10 cycles/2 mins															
120	100	100	80	80	80	80	80	80	80	80	80	80	80	80	80	80	ETT Drug Administration																	
160	160	160	120	120	120	120	120	120	120	120	100	100	100	100	100	100	Adrenaline	10 time the IV dose																
																	Atropine	30 mcg/kg																
																	All other drugs double the IV dose																	
																	Dilution For ETT Drug Admin																	
																	Newborn/Infants.....	1 ml																
																	Small Child	3 - 5 mls																
																	Large Child	5 - 10 mls																
																	Suction Catheter Size For ETT																	
																	6 FG	3 mm																
																	8 FG	3.5 - 5.5 mm																
																	10 FG	6 mm																
																	CO2 Retentions Signs/Symptoms																	
																	• Pupillary Dilatation	• Sweating																
																	• CNS Depression	• Hypertension																
																	• Bounding Pulse	• Tachycardia																
																	Hypoxia Signs/Symptoms																	
																	• Pallor	• Restlessness																
																	• Bradycardia	• Cyanosis																
																	• Hypotension	• Tachypnea																
																	• Apnoea	• Tachycardia																
																	• Lethargy	• Bradycardia (late)																
																	Neb Adrenaline for Croup Tx Indications																	
																	• Signs of Hypoxia (as above)																	
																	• SPO2 of <92 on air																	
																	• Marked use of Resp Accessory Muscles																	
																	Meningococcal Tx Indications																	
																	• Purpuric Rash And	• Headache	• Tachycardia															
																		• Fever	• Hypotension															
																	TCA Overdose Tx Indications																	
																	• QRS >0.12 (>0.16 severe toxicity)																	
																	• QT prolongation (>1/2 R-R interval)																	
																	• Sinus Tachycardia																	
																	Intraosseous Needle Contraindications																	
																	• Limb traumatised or infected																	
																	• Site cannot be adequately cleaned																	
																	Anaphylaxis Tx Indications																	
																	• Resp Distress with Bronchospams OR																	
																	• Less than adequate perfusion OR																	
																	• Altered Conscious State With Either	• Angionema																
																		• Urticaria																
																		• GIT Manifestations																
																	GHB Overdose																	
																	Tx Indications																	
																	• Agitated and Aggressive																	
																	Aim Of Tx																	
																	• Achieve arousable drowsiness																	
																	Sedate To Intubate																	
																	INDICATIONS GCS <10 &																	
																	• Respiratory Failure																	
																	• Status Epilepticus																	
																	CONTRAINDICATIONS																	
																	• Head injury & strong jaw tone																	
																	• Potential impossible intubation																	
																	• No functional electronic capnograph																	
																	APGAR																	
																	Appearance	0	1	2														
																	Blue/Pale	Body Pink	All Pink	All Pink														
																	Pulse	Absent	<100	>100														
																	Grimace	None	Grimace	Cries														
																	Activity	Limp	Extremities	Active														
																	Respirations	Absent	Slow/Weak	Good crying														
																	Adequate Perfusion																	
																	Age	Pulse		Systolic BP														
																	Newborn	120 - 160		N/A														
																	Infant	100 - 160		> 70 mmhg														
																	Sm Child	80 - 120		> 80 mmhg														
																	Lg Child	80 - 120		> 90 mmhg														
																	Date Of Card: 12 July 2006																	
																	This card has been prepared for educational & training purposes only. It is a guide only for use with Victorian Ambulance Guidelines April 2006 & must only be used by those properly trained in it's use. The user accepts all responsibility in it's use.																	
																	0.3	0.5	0.7	1	1.2	1.4	1.6	1.8	2	2.2	2.4	2.6	3.3	3.6	4	4.3	4.6	
																		0.7	1	1.4	2	2.4	2.8	3.2	3.6	4	4.4	4.8	5.2	6.6	7.2	8	8.6	9.2

Cut along this line before laminating

Fold Line

Fold Line

Cut along this line before laminating

Cut along this line before laminating